

## A Ten-Year Investigation on Ozone and Its Precursors at Kemaman, Terengganu, Malaysia

Marzuki Ismail<sup>a</sup>, Samsuri Abdullah<sup>a</sup>, Fong Si Yuen<sup>a</sup> and Nurul Adyani Ghazali<sup>b</sup>

<sup>a</sup> School of Marine Science and Environment, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu, Malaysia

<sup>b</sup> School of Ocean Engineering, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu, Malaysia

### Abstract

10 years of continuous monitoring data (2000-2010) from Air Quality Division, Malaysian Department of Environment are used to investigate the relationships between ambient levels of ozone (O<sub>3</sub>), nitric oxide (NO) and nitrogen dioxide (NO<sub>2</sub>) as a function of NO<sub>x</sub> in Kemaman, Malaysia, wherewith the dominant sources of ozone precursors are industrial activities and road traffic. In addition, variation of oxidant OX (O<sub>3</sub> and NO<sub>2</sub>) concentration with NO<sub>x</sub> was also examined. The analyses established a weak linear relationship between O<sub>3</sub> and NO<sub>x</sub>, signifying that Kemaman area is not NO<sub>x</sub> sensitive but possibly VOC-sensitive area. The level of [OX] is influenced by NO<sub>x</sub>-independent and NO<sub>x</sub>-dependent contributions. The former is due to regional background O<sub>3</sub> concentration while the latter correlates to the local level of primary pollution. The measured concentrations of the pollutants varied as a function of time. The analyses also show that the diurnal cycle of ground level ozone concentration has a mid-day peak while lower concentration occurs at night time.

**Keywords:** ground level ozone; solar radiation intensity; diurnal cycle; oxidant; Kemaman

### 1. Introduction

Ozone is a secondary pollutant resulting from photochemical reaction of variety of natural and anthropogenic precursors; mainly volatile organic compound (VOC) and oxides of nitrogen (NO<sub>x</sub>). The formation of ground level ozone depends on the intensity of solar radiation, the absolute concentration of NO<sub>x</sub> and VOCs, and the ratio of NO<sub>x</sub> to VOCs. Ground level ozone comes from two major sources namely regional background ozone primarily from the influx of ozone from the stratosphere and in situ local ozone production mainly via precursors, such as NO<sub>x</sub> and VOC, emitted from variety of anthropogenic and natural sources (Clapp and Jenkin, 2001). Ozone at high concentrations could induce several adverse effects on human health and the environment (Kampa and Castanas, 2007). The concentration of photochemical oxidant (O<sub>3</sub> and NO<sub>2</sub>) can be decreased by controlling their precursors; Nitrogen Oxides, NO<sub>x</sub> (NO and NO<sub>2</sub>) and VOCs (Geng *et al.*, 2007). However, the efficiency of emission control also depends on the relationship between

primary and secondary pollutants, as well as ambient meteorological factors. Owing to the chemical coupling of O<sub>3</sub> and NO<sub>x</sub>, the levels of O<sub>3</sub> and NO<sub>2</sub> are inextricably linked. The changes in the local level of O<sub>3</sub> and NO<sub>2</sub> will lead to an increasing background level. In this study, ten years ambient concentration data of O<sub>3</sub>, NO, NO<sub>2</sub>, and NO<sub>x</sub> were used for the investigation of the hourly variation of NO, NO<sub>2</sub>, NO<sub>x</sub> and O<sub>3</sub>. The variation of oxidant OX (O<sub>3</sub> and NO<sub>2</sub>) concentration with NO<sub>x</sub> was also investigated, contributing to a better understanding of the atmospheric sources of OX at this Kemaman area.

### 2. Materials and Methods

#### 2.1. Site description

Kemaman (4°12' N, 103°18' E) is situated in the Terengganu State, a developing Malaysian town flanked by Kerteh Petrochemical Industrial Area at the north and the industrializing and urbanizing Gebeng Industrial Area at the south (Fig. 1). In this area, there are dominant sources of ozone precursors related to industrial activities and road traffic.



Figure 1. Location of air quality monitoring station in Kemaman

## 2.2. Source of data

In this study, the concentration of  $O_3$ , NO,  $NO_2$ , and  $NO_x$  data consisting of 120 months from January 2000 to December 2010 (excluding year 2008 due to the unavailability of the data for that particular year) acquired from the Air Quality Division, Malaysian Department of Environment for Sekolah Rendah Bukit Kuang Station (Station ID: CA 0002) located in Kemaman District, one of the earliest operational stations in Malaysia. The monitoring network was installed, operated and maintained by Alam Sekitar Malaysia Sdn. Bhd (ASMA) under concession by the Department of Environment Malaysia (Afroz *et al.*, 2003). Tropospheric ozone concentrations data was recorded by using a system based on the Beer-Lambert law for measuring low ranges of ozone in ambient air manufactured by Teledyne Technologies Incorporated (Model 400E). A 254 nm UV light signal is passed through the sample cell where it is absorbed in proportion to the amount of ozone present. Every three seconds, a switching valve alternates measurement between the sample stream and a sample that has been scrubbed of ozone. The result is a true, stable

ozone measurement. The concentration of ozone precursors,  $NO_x$  (NO and  $NO_2$ ) were determined using the chemiluminescence measurement principle, couple with state-of-the-art microprocessor technology for monitoring high and medium levels of nitrogen oxides (Teledyne Models 200A). For quality control and quality assurance of the air monitoring data, all monitoring instruments were calibrated regularly by ASMA. For gas monitoring such as ozone and nitrogen oxides, the instruments were scheduled to have daily auto calibration using zero air (clean ambient air, free from contaminant) and standard gas concentration. The instruments have been calibrated manually using its individual calibration gas every two weeks. The hourly data was also checked for validation before it can be transferred to the Department of Environment.

## 2.3. Data analysis

In order to establish the hourly trend of selected pollutants, boxplot was performed by using Microsoft Excel Spreadsheet 2013. In descriptive statistics, a boxplot is a convenient way of graphically depicting groups of numerical data through their quartiles.




Figure 2. Hourly variation of mean values of NO, NO<sub>2</sub>, NO<sub>x</sub>, and O<sub>3</sub> concentrations from 2000-2010\* (excluding year 2008)

Boxplot is a graphic representation of a distribution by a rectangle, the ends of which mark the maximum and minimum values, and in which the median and first and third quartiles are marked by lines parallel to the ends. The individual box plot is a visual aid to examining key statistical properties of a variable such as median, mean and quartiles. Analyses of variance and correlation analysis (the correlation is significant at the 0.05 level (2-tailed)) were conducted using STATGRAPHICS® Centurion XVI.I statistical software package.

### 3. Results and Discussion

#### 3.1. Hourly variation of O<sub>3</sub>, NO, NO<sub>2</sub> and NO<sub>x</sub> concentrations

The 10 years (2000-2010, excluding year 2008) average diurnal variation of NO, NO<sub>2</sub>, NO<sub>x</sub> and O<sub>3</sub> concentrations were shown in Fig. 2.

Based on Fig. 3, the diurnal cycle of ozone concentration reaches a peak during the middle of the day, while there was a lower ozone concentration during nighttime (Ramli *et al.*, 2010). This is due to the photochemical reaction of O<sub>3</sub> precursors, such as volatile organic compound (VOC), with ambient air from natural sources and the long distance transport of NO<sub>x</sub> (Pires *et al.*, 2012). The ozone concentration slowly increases after the sun rises until it reached its maximum during the daytime and then slowly decreases until the next morning. From 08:00 to 13:00-14:00 hours, an increase in solar radiation results in the increasing of ozone concentration (Ulke and Maezzo, 1998). The average data of ozone concentration and solar radiation shows that the ozone concentration is highly correlated to solar radiation (J/m<sup>2</sup>.hr). It was found that the correlation coefficient between ozone and solar radiation was 0.67 from the statistical analysis, at the 0.05 level (2-tailed).




Figure 3. Hourly variation of mean values of solar radiation from 2000-2010\* (excluding 2008)

The diurnal cycle of NO, NO<sub>2</sub> and NO<sub>x</sub> are shaped like double waves. The morning peak of NO<sub>2</sub> occurs about two hours after the morning peak of NO (08:00). The O<sub>3</sub> peak occurs at 14:00 (Azmi *et al.*, 2010). It appears about 6 to 8 hours after the NO (08:00) and NO<sub>2</sub> (10:00) peak. After the morning peak (08:00), NO diminishes until it reaches lower value at (13:00-14:00). It was found that, both NO and NO<sub>2</sub> decreases correlate with an increase in O<sub>3</sub> (Ramli *et al.*, 2010). The increasing of NO<sub>2</sub> has caused the decreasing of O<sub>3</sub> concentrations (Ghazali *et al.*, 2008). NO is a primary contaminant, whereas O<sub>3</sub> and a large percentage of NO<sub>2</sub> are secondary contaminants, formed through a set of complex reaction. NO is converted to NO<sub>2</sub> via a reaction with O<sub>3</sub> as in Eq. (1). During daytime, NO<sub>2</sub> is converted back to NO as a result of photolysis, leads to regeneration of O<sub>3</sub> as in Eq. (2). The vertical mixing layer of also influenced the air pollution concentrations (Banan *et al.*, 2013). During the day, pollutants would be diluted when mixing layer rises and be limited to inside nocturnal planetary boundary layer (NPBL) during nighttime. Emitted pollutants such as NO and NO<sub>2</sub> are kept beneath this inversion, which may cause the hourly NO<sub>x</sub> concentration to increase during the night (Han *et al.*, 2011)

### 3.2. Chemical coupling of O<sub>3</sub>, NO and NO<sub>2</sub>

The inter conversion of O<sub>3</sub>, NO and NO<sub>2</sub> under atmospheric conditions is generally dominated by the following reactions (Kenty *et al.*, 2007)



These equations form a cycle with no net chemistry; the overall effect of reaction Eq. (2) is the reverse of reaction Eq. (1). These reactions therefore represent a closed system which has the overall effect of partitioning NO<sub>x</sub> between its component forms of NO and NO<sub>2</sub>, and oxidant (OX) between its component forms of O<sub>3</sub> and NO<sub>2</sub>, but leaving a total mixing ratio of both NO<sub>x</sub> and OX unchanged. During daytime hours, NO, NO<sub>2</sub> and O<sub>3</sub> are typically equilibrated on the timescale of a few minutes, a condition usually referred to as photo stationary state. NO, NO<sub>2</sub> and O<sub>3</sub> concentrations are related by the expression [NO]. [O<sub>3</sub>]/[NO<sub>2</sub>] = J<sub>2</sub>/k<sub>1</sub>, where J<sub>2</sub> is the rate of NO<sub>2</sub> photolysis and k<sub>1</sub> is the rate coefficient for the reaction of NO with O<sub>3</sub> (Han *et al.*, 2011).

The variation of the mean value of J<sub>2</sub>/k<sub>1</sub> over time, obtained using 10 years data at Kemaman of NO, NO<sub>2</sub> and O<sub>3</sub> is shown in Fig. 4. The mean values of J<sub>2</sub>/k<sub>1</sub> vary between 3.225 and 12.869 ppb and the maximum value occurred at 15:00. This provides adequate description of diurnal averaged observations, consistent with the chemical coupling of these species being dominated by reactions Eq. (1) and Eq. (2).

Fig.5 shows the variation in daytime O<sub>3</sub> concentration as a function of the NO<sub>2</sub>/NO ratio. It was found that the level of O<sub>3</sub> increases with an increase in NO<sub>2</sub>/NO ratio. O<sub>3</sub> increases rapidly with a small value of NO<sub>2</sub>/NO ratio, this may be implicated that when O<sub>3</sub> was at low levels, the reactions of production of O<sub>3</sub> was dominated reaction (Han *et al.*, 2011). When O<sub>3</sub> reached at about 35 ppb, it tended to remains relatively stable. This shows that at higher levels, O<sub>3</sub> is close to reaching a photo stationary state. The output shows the results of fitting a logarithmic-X model to describe




Figure 4. Variation of mean values of  $J_2/k_1$

the relationship between  $O_3$  concentrations and  $NO_2/NO$ . The equation of the fitted model is  $O_3$  (ppb) =  $13.6992 + 12.3397 \cdot \ln(NO_2/NO)$ . Since the result of  $P$ -value from the ANOVA is less than  $0.05$ , there is a statistically significant relationship between  $O_3$  and  $NO_2/NO$  at the 95.0% confidence level. The logarithmic function can be used to forecast the  $[O_3]$  during the daytime. The correlation coefficient equals to  $0.70$ , indicating a moderately strong relationship between the variables.

### 3.3. Diurnal variation of $[OX]$

Fig. 6 shows the variation in the mean value of  $OX$  concentration.  $OX$  concentration shows a mid-day peak and lower nighttime concentrations which is similar to the variation of  $O_3$  concentration. The  $OX$  concentration slowly raises after the sunrise, reaches a maximum

during the day and then decreases until the next morning. This is due to the photochemical  $O_3$  formation (Pires *et al.*, 2012). Fig. 7 shows the variation of  $[NO_2]/[OX]$ . A smaller ratio of  $[NO_2]/[OX]$  is due to the higher concentration of  $O_3$  during the day. This difference in the partitioning of  $NO_2$  and  $O_3$  may be related to the rate of chemical processes or the time available for them to occur. For instance, the higher ratio of  $[NO_2]/[OX]$  was due to the lower  $O_3$  concentration during the day.

### 3.4. Local and regional contributions to oxidant

The variation of  $OX$  concentration against the level of  $NO_x$  is shown in Fig. 8. The total  $OX$  appears to increase linearly with  $NO_x$  over the entire range considered, such that the level of  $OX$  at Kemaman has  $NO_x$ -independent contribution and  $NO_x$ -dependent contribution. The  $NO_x$ -independent contribution can




Figure 5. Relationship of  $O_3$  concentration with the  $[NO_2]/[NO]$  ratio from 2000-2010\* (excluding year 2008)




Figure 6. Variation of mean values of OX




Figure 7. Variation of mean values of NO<sub>2</sub>/OX ratio

be considered as a regional contribution which equates to the regional background of O<sub>3</sub> level, while the NO<sub>x</sub>-dependent contribution can be considered as local contribution and is correlated with the level of primary pollution. The result of this study is similar as performed by Clapp and Jenkin (2001).

The local OX source has probable contributions from direct NO<sub>2</sub> emission, thermal reaction of NO with O<sub>2</sub> at high NO<sub>x</sub> and common source emission of species which promote NO to NO<sub>2</sub> conversion. The regional contribution to O<sub>3</sub> concentration was known to be at about 18.22 ppb according to the intercept of the relationship between OX concentration and NO<sub>x</sub> concentration.

### 3.5. Correlations between O<sub>3</sub> and NO<sub>x</sub>

Fig. 9 shows that the concentration of O<sub>3</sub> and NO<sub>x</sub> were linearly correlated. The correlation coefficient equals to  $-0.165$ , indicating a relatively weak relationship between the variables (O<sub>3</sub> and NO<sub>x</sub>). A weak correlation between O<sub>3</sub> and NO<sub>x</sub> indicates that Kemaman area is probably not NO<sub>x</sub>-sensitive but possibly VOC-sensitive. This inverse relationship of O<sub>3</sub> and NO<sub>x</sub> is due to the titration of O<sub>3</sub> during the daytime (Banan *et al.*, 2013; Nishanth *et al.*, 2014). Based on Fig. 10, the NO<sub>2</sub>/NO<sub>x</sub> ratio decreased greatly as NO<sub>x</sub> increased that could be newly emitted could also support the VOC-sensitive nature in Kemaman. This finding is corresponding as




Figure 8. Relationship between [OX] with the level of [NO ] from 2000-2010\* (excluding year 2008)






Figure 9. Correlations between  $[O_3]$  and  $[NO_x]$  from 2000-2010\* (excluding year 2008)

Figure 10. Relationship  $NO_2/NO_x$  ratio from 2000-2010\* (excluding year 2008)

found by Song *et al.* (2011). VOC was emitted from the industrialized area and may also be emitted by traffic, and more detailed relationship among  $NO_x$ , VOC and  $O_3$  in this area need to be explored through future studies.

#### 4. Conclusions

This study analyses the concentrations of  $NO$ ,  $NO_2$ ,  $NO_x$  and  $O_3$  for 10 years from 2000 to 2010 (excluding year 2008). The results indicate that the diurnal cycle of ozone concentration has a mid-day peak (14:00) and lower nighttime concentrations. The ozone concentration slowly rises after the sun rises (08:00), reaching a maximum during daytime and then decreases until the next morning. This is due to the photochemical of ozone formation. In this study, it was also found a weak linear relationship between  $O_3$  and  $NO_x$ , suggesting that this Kemaman area is not  $NO_x$  sensitive but possibly VOC-sensitive area. A logarithmic relationship between  $O_3$  concentration and  $[NO_2]/[NO]$  ratio was also found which could be useful in  $[O_3]$  forecasting and air pollution control strategies. The level of  $[OX]$  is influenced by  $NO_2$ -independent and  $NO_2$ -dependent contributions. The former is due to regional background  $O_3$  concentration and the latter correlates to the local level of primary pollution. The regional background  $O_3$  concentration in Kemaman is about 18.22 ppb

#### Acknowledgements

This study was funded by Fundamental Research Grant Scheme (FRGS) FRGS/2/2013/STWN01/UMT/02/1 (VOT 59312) and Universiti Malaysia Terengganu Scholarship Scheme (BUMT). The authors would also like to thank the Air Quality Division, Malaysian Department of Environment (DOE) for the air quality data.

#### References

- Afroz R, Hassan MN, Ibrahim NA. Review of air pollution and health impacts in Malaysia. *Environmental Research* 2003; 92(2): 71-77.
- Azmi SZ, Latif MT, Ismail AS, Juneng L, Jemain AA. Trend and status of air quality at three different monitoring stations in the Klang Valley, Malaysia. *Air Quality, Atmosphere and Health* 2010; 3(1): 53-64.
- Banan N, Latif MT, Juneng L, Ahamad F. Characteristics of surface ozone concentrations at stations with different backgrounds in the Malaysian Peninsula. *Aerosol and Air Quality Research* 2013; 13(3): 1090-106.
- Clapp LJ, Jenkin ME. Analysis of the relationship between ambient levels of  $O_3$ ,  $NO_2$ , and  $NO$  as a function of  $NO_x$  in the UK. *Atmospheric Environment* 2001; 35(36): 6391-405.
- Geng F, Zhao C, Tang X, Lu G, Tie X. Analysis of ozone and  $VOC_s$  measured in Shanghai: A case study. *Atmospheric Environment* 2007; 41(5): 989-1001.
- Ghazali NA, Ramli NA, Yahaya AS, Yusof NF, Sansuddin N, Madhoun WA. Temporal analysis and prediction of ozone concentrations in Gombak, Selangor. *International Conference on Construction and Building Technology* 2008; 221-32.
- Han S, Bian H, Feng Y, Liu A, Li X, Zeng F, Zhang X. Analysis of the relationship between  $O_3$ ,  $NO$  and  $NO_2$  in Tianjin, China. *Aerosol and Air Quality Research* 2011; 11: 128-39.
- Kampa M, Castanas E. Human health effects of air pollution. *Environmental Pollution* 2007; 151(2): 362-67.
- Kenty KL, Poor ND, Kronmiller KG, McClenny W, King C, Atkeson T, Campbell SW. Application of CALINE4 to roadside  $NO/NO_2$  transformations. *Atmospheric Environment* 2007; 41(20): 4270-80.
- Nishanth T, Praseed KM, Kumar MKS, Valsaraj KT. Observational study of surface  $O_3$ ,  $NO_x$ ,  $CH_4$  and total NMHCs at Kannur, India. *Aerosol and Air Quality Research* 2014; 14: 1074-88.

- Pires JCM, Alvim-Ferraz MCM, Martins FG. Surface ozone behaviour at rural sites in Portugal. *Atmospheric Research* 2012; 104-105: 164-71.
- Ramli NA, Ghazali NA, Yahaya AS. Diurnal fluctuations of ozone concentrations and its precursors and prediction of ozone using multiple linear regressions. *Malaysian Journal of Environmental Management* 2010; 11(2): 57-69.
- Song F, Shin JY, Jusino-Atresino R, Gao Y. Relationship among the springtime ground-level  $\text{NO}_x$ ,  $\text{O}_3$  and  $\text{NO}_3$  in the vicinity of highways in the US East Coast. *Atmospheric Pollution Research* 2011; 2(3): 374-83.
- Ulke AG, Mazzeo NA. Climatological aspects of the daytime mixing height in Buenos Aires City, Argentina. *Atmospheric Environment* 1998; 32(9): 1615-22.
- 

*Received 21 April 2015*

*Accepted 8 June 2015*

**Correspondence to**

Dr. Marzuki Ismail  
School of Marine Science and Environment,  
Universiti Malaysia Terengganu  
21030 Kuala Terengganu,  
Terengganu,  
Malaysia  
Tel: +60 9668 3548  
Fax: +60 9669 4669  
E mail: marzuki@umt.edu.my