

A SURVEY ON HABITAT-SPECIFIC COMPOSITION OF THE
BATS AT INSTITUTE OF INDUSTRIAL CAMPUS AREA

JULIAH BINTI YAHYA

EKARIS SAHARAH TENGKOLAH

KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MELAKA

2005

Perustakaan

Kolej Universiti Sains Dan Teknologi Malaysia (KUSTEM)

1100036883

LP 42 FST 4 2005



1100036883

A survey on ectoparasites composition of the bats at vicinity of KUSTEM campus area / Zuliha Yahya.



PERPUSTAKAAN

**KOLEJ UNIVERSITI SAINS & TEKNOLOGI MALAYSIA
21030 KUALA TERENGGANU**

1100036883

1100036883

Lihat sebelah

HAK MILIK
PERPUSTAKAAN KUSTEM

A SURVEY ON ECTOPARASITES COMPOSITION OF THE BATS AT
VICINITY OF KUSTEM CAMPUS AREA

By

Zuliha binti Yahya

Research Report submitted in partial fulfillment of
the requirements for the degree of
Bachelor of Applied Science (Biodiversity Conservation and Management)

Department of Biological Sciences
Faculty of Science and Technology
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MALAYSIA
2005

This project should be cited as:

Zuliha, Y. 2005. A survey on ectoparasites composition at vicinity of KUSTEM campus area. Undergraduate thesis, Bachelor of Applied Science in Biodiversity Conservation and Management, Faculty of Science and Technology, Kolej Universiti Sains dan Teknologi Malaysia, Terengganu. 70p.

No part of this project report may be produced by any mechanical, photographic, or electronic process, or in the form of phonographic recording, nor may it be stored in a retrieval system, transmitted, or otherwise copied for public or private use, without written permission from the author and the supervisor(s) of the project.



JABATAN SAINS BIOLOGI
FAKULTI SAINS DAN TEKNOLOGI
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MALAYSIA

PENGAKUAN DAN PENGESAHAN LAPORAN
PROJEK PENYELIDIKAN I DAN II

Adalah ini diakui dan disahkan bahawa laporan penyelidikan bertajuk: .

Survey Ke Atas Komposisi Ektoparasit Pada Kelawar Di Persekutaran Kawasan KUSTEM oleh Zuliha binti Yahya, no. matrik: UK7182 telah diperiksa dan semua pembetulan yang disarankan telah dilakukan. Laporan ini dikemukakan kepada Jabatan Sains Biologi sebagai memenuhi sebahagian daripada keperluan memperolehi Ijazah ... Bachelor Sains Gunaan (Pemuliharaan dan Pengurusan Biodiversiti), Fakulti Sains dan Teknologi, Kolej Universiti Sains dan Teknologi Malaysia.

Disahkan oleh:

Penyelia Utama

Nama: **WONG CHEE HO**
Cop Rasmi: Pensyarah
Cop Rasmi: Jabatan Sains Biologi
Fakulti Sains dan Teknologi
Kolej Universiti Sains dan Teknologi Malaysia
(KUSTEM)
21030 Kuala Terengganu.

Tarikh:

Penyelia Kedua (jika ada)

Nama:

Cop Rasmi

Tarikh:

Ketua Jabatan Sains Biologi

Nama: **PROF. MADYA DR. MAYSAM MAT AMIN**
Cop Rasmi: **PROF. MADYA DR. MAYSAM MAT AMIN**
Cop Rasmi: Jabatan Sains Biologi
Fakulti Sains dan Teknologi
Kolej Universiti Sains dan Teknologi Malaysia
(KUSTEM)
21030 Kuala Terengganu
21030 Kuala Terengganu

Tarikh:

ACKNOWLEDGEMENT

First of all, thank God for giving me this chance and abilities to fulfill my report research. I would like to acknowledge the contribution of a number of people, who have volunteered many hours of their time in assisting to finish my work in preparing this report. Special thanks to my supervisor Mr. Wong Chee Ho for his suggestions and idea in preparing, arranging and completing my report.

The establishment of this report in such a timely manner would not have been possible without the support and generosity of Mr.Amir, the lecturer of Biological Science Department. Sincerely appreciation to him for his willingness to help me completed this project. His suggestions and guidance were greatly appreciated.

Many of my family members offered prayers and provided materials and suggestions for this report as well. I am deeply value for their contribution. I will always remember the reason for their contribution and let glory and honor to them.

During the field work, I was assisted by lab assistant Mr. Mohamad Embong. A lot of assistance was given by this kind man, including in setting up net in sampling site. He also gave me much helpful in providing materials in identifying work and guiding me in lab work.

Last but not least I would like to thank my friends, Noor Azlin Asnam, Wan Daraputri Razali and Widad Fadhlullah for their assistance during the field work and their suggestions and comments were very helpful for me to complete this report.

TABLE OF CONTENTS

CONTENT	Page
ACKNOWLEDGEMENT	i
TABLE OF CONTENTS	iii
LIST OF TABLES	vii
LIST OF FIGURES	viii
LIST OF APPENDICES	ix
ABSTRACT	x
ABSTRAK	xi
1.0 INTRODUCTION	1
OBJECTIVES	2
2.0 LITERATURE REVIEW	3
2.1 Parasitism	3
2.2 Theory of parasitism	4
2.3 Classifications of ectoparasites	5
2.3.1 Endoparasites	6
2.3.2 Ectoparasites	6
2.4 Types of ectoparasites	7
2.4.1 Mites	7
2.4.2 Fleas	9

2.4.3	Bugs	9
2.4.4	Flies	11
2.4.5	Louses	12
2.5	Adaptations of ectoparasite's morphology	13
2.5.1	Mites	13
2.5.2	Fleas	14
2.5.3	Bugs	14
2.5.4	Louses	14
2.6	Harmful of ectoparasites	15
2.6.1	Harmful to bats	15
2.6.2	Harmful to humans	15
2.7	Bats	16
2.7.1	Megachiroptera	17
2.7.2	Microchiroptera	17
2.8	The parasites of bats and lifecycle	18
2.8.1	Fully on bats	20
2.8.2	In the immediate environment of bats	20
2.8.3	Contact the bats only just to feed	20
2.9	Relationships	21
2.9.1	Host-parasite interrelationship	21
2.9.2	Ectoparasites-bats relationship	21

3.0	METHODOLOGY	
3.1	Sampling site	23
3.2	Material	25
3.3	Removal bats from nets	25
3.4	Lab analysis	26
3.5	Slide observations	26
3.6	Identifying	27
3.6.1	Mite	27
3.6.2	Flea	28
3.6.3	Bug	28
3.6.4	Fly	28
3.6.5	Lice	28
3.7	Data analysis	29
3.7.1	Relative abundance	29
3.7.2	Species abundance	29
3.7.3	Shannon Weiner Index	29
3.7.4	Simpson Index	30
3.7.5	Infected Index	30
3.7.6	Prevalence Index	30
4.0	RESULTS	
4.1	Bats abundance and diversity in general	31
4.2	Ectoparasites	37

4.2.1	Infected index of number of fleas for examined bats	41
4.2.2	Prevalence index for different gender of bats	44
5.0	DISCUSSION	55
5.1	Bats	55
5.2	Ectoparasites	59
6.0	CONCLUSION	61
REFERENCES		62
APPENDIX		67

LIST OF TABLES

TABLE		Page
4.1	Percentage of relative abundance according to the species captured	33
4.2	Value of Shannon-Weiner and Simpson Diversity Indices	35
4.3	Percentage of infected bats	36
4.4	Percentage of different types of ectoparasites on two species of bats	41
4.5	Infected index of fleas for examined bats	42
4.6	Prevalence index for different gender of bats	44

LIST OF FIGURES

FIGURES	Page
3.1 Sampling site at Kg.Padang Nenas, Terengganu	24
3.2 Another sampling site at Kg. Padang Nenas, Terengganu	24
4.1 Graph shows the number of captured bats throughout four months of sampling.	32
4.2 Graph shows the number of ectoparasites for four months	37
4.3 Graph shows the different ectoparasites on two species of bats	39
4.4 Graph shows the numbers of ectoparasites for different gender of bats	46
4.5.1 Body parts of <i>Nycteribia sp.a</i> ; abdomen	48
4.5.2 Another parts of <i>Nycteribiasp</i>	48
4.5.3 Legs and posterior parts of <i>Nycteribia sp</i> ; <i>lab</i> ;labium, <i>m</i> ;mentum	49
4.5.4 Leg's structure of <i>Nycteribia sp.t</i> ; <i>tarsus,mt</i> ; metatarsus, <i>u</i> :ungues or claw	49
4.5.5 An anterior parts of a <i>Nycteribia sp.f</i> ; femur, <i>h</i> ; head.	50
4.5.6 The posterior parts of a <i>Nycteribia sp.t</i> ; <i>tibia,lab</i> ; labium with teeth	50
4.6 <i>Ctenocephalides sp a</i> ;abdomen, <i>f</i> ;femur, <i>mp</i> ;palp of first maxilla, <i>ta</i> ;tarsus with claws.	51
4.7 Analysis indices value	53

LIST OF APPENDIX

Appendix	Page
I The list of individual for each species recorded for the whole sampling period.	67

A survey On Ectoparasites Composition Of The Bats At Vicinity Of KUSTEM Campus Area.

ABSTRACT

A study was conducted on the ectoparasites composition of bats at vicinity of KUSTEM campus area. The objectives of this study including examining and comparing the ectoparasites among different species of bats. Sampling sites have been chosen randomly for the placement of mist nets. Mist nets were used to trap the bats. The sampling has been conducted for five days monthly from October 2004 until January 2005 respectively. Overall, 67 individuals of *Cynopterus brachyotis* and 12 individuals of *Cynopterus horsfieldii* were caught during the four months of sampling. From the 79 individuals of the fruit bats that have been recorded, there were 47 individuals of ectoparasites comprised by 36 individuals of *Nycteribia sp* and 11 individuals of *Ctenocephalides sp* were collected. *Cynopterus brachyotis* showed the higher percentage of infected bats while *Cynopterus horsfieldii* showed the higher prevalence value. *Nyteribia sp* composition was commonly found on *Cynopterus brachyotis* while *Ctenocephalides sp* composition was on *Cynopterus horsfieldii*. The Shannon-Weiner index value is 0.5442 while Simpson index value is 0.3585. In conclusions of the present study indicate that ectoparasites found on bats influenced by the gender and of bats and also the seasons.

Survey Ke Atas Komposisi Ektoparasit pada Kelawar di Persekutaran Kawasan KUSTEM

ABSTRAK

Kajian ini telah dijalankan untuk mengkaji komposisi ektoparasit pada kelawar di persekitaran kawasan KUSTEM. Objektif kajian ini termasuklah mengkaji dan membandingkan ektoparasit-ektoparasit yang terdapat pada kelawar yang berlainan spesies. Kawasan persampelan telah dipilih secara rawak untuk pemasangan jaring kabus. Jaring kabus telah digunakan untuk menangkap kelawar. Persampelan ini telah dijalankan selama 5 hari pada setiap bulan dari Oktober 2004 hingga Januari 2005 masing-masing. Secara keseluruhannya, 67 individu *Cynopterus brachyotis* dan 12 individu *Cynopterus horsfieldii* telah ditangkap selama empat bulan persampelan. Daripada 79 individu kelawar buah ini, 47 ektoparasit yang terdiri daripada 36 individu *Nycteribia sp* dan 11 individu *Ctenocephalides sp* telah dikumpulkan. *Cynopterus brachyotis* menunjukkan nilai peratusan kelawar yang dijangkiti lebih tinggi manakala *Cynopterus horsfieldii* menunjukkan nilai kelaziman yang lebih tinggi. Komposisi *Nycteribia sp* biasanya adalah pada *Cynopterus brachyotis* manakala komposisi *Ctenocephalides sp* adalah pada *Cynopterus horsfieldii*. Nilai index Shannon-Weiner ialah 0.5442 manakala nilai bagi index Simpson ialah 0.3585. Kesimpulannya, kajian ini menunjukkan bahawa ektoparasit yang dijumpai pada kelawar dipengaruhi oleh jantina dan habitat kelawar dan juga musim.