

GUARIBA PITCH (FICUS DELTOIDEA) JACK.
SEEDLING GROWTH IN NATURAL AND ARTIFICIAL HABITAT
AND ARTIFICIAL SEEDBED

NURLIYANA BINTI ZAKI

FAKULTI SAINS DAN TEKNOLOGI

UNIVERSITI SAINS SABAH, SABAH, MALAYSIA

2006

LP
44
FST
5
2006

CH: 4806

Perpustakaan Universiti Malaysia Terengganu (UMT)
1100046118



LP 44 FST 5 2006



1100046118

Comparison of (*Ficus Deltoidea*) Jack. seedlings growth between natural habitat and artificial medium / Nurliyana Ab. Razak.

PERPUSTAKAAN

**KOLEJ UNIVERSITI SAINS & TEKNOLOGI MALAYSIA
21030 KUALA TERENGGANU**

1100046118

Lihat sebelah

HAK MILIK
PERPUSTAKAAN KUSTEM

**COMPARISON OF *FICUS DELTOIDEA* JACK. SEEDLINGS GROWTH
BETWEEN NATURAL HABITAT AND ARTIFICIAL MEDIUM**

By
Nurliyana bt Ab Razak

Research Report submitted in partial fulfillment of
the requirement for the degree of
Bachelor of Applied Science (Biodiversity Conservation and Management)

Department of Biological Sciences
Faculty of Science and Technology
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MALAYSIA
2006

This project should be cited as:

Nurliyana, A. R. 2006. Comparison of *Ficus deltoidea* Jack. seedlings growth between natural habitat and artificial medium, Undergraduate thesis, Bachelor of Applied Science in Biodiversity Conservation and Management, Faculty of Science and Technology, Kolej Universiti Sains Dan Teknologi Malaysia. 28p.

No part of this project report may be produced by any mechanical, photographic, or electronic process, or in the form of phonographic recording, nor may it be stored in a retrieval system, transmitted, or otherwise copied for public or private use, without written permission from the author and the supervisors of the project.

1100046118



JABATAN SAINS BIOLOGI
FAKULTI SAINS DAN TEKNOLOGI
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MALAYSIA

PENGAKUAN DAN PENGESAHAN LAPORAN
PROJEK PENYELIDIKAN I DAN II

Adalah ini diakui dan disahkan bahawa laporan penyelidikan bertajuk: COMPARISON OF FICUS DELTOIDEA JACK. SEEDLINGS GROWTH BETWEEN NATURAL HABITAT AND ARTIFICIAL MEDIUM oleh Nurliyana Ab. Razak no. matrik: UK 8873 telah diperiksa dan semua pembetulan yang disarankan telah dilakukan. Laporan ini dikemukakan kepada Jabatan Sains Biologi sebagai memenuhi sebahagian daripada keperluan memperolehi Ijazah Sarjana Muda Sains Gunaan (Pemuliharaan dan Pengurusan Biodiversiti) Fakulti Sains dan Teknologi, Kolej Universiti Sains dan Teknologi Malaysia.

Disahkan oleh:

Penyelia Utama **JAMILAH MOHD SALIM @ HALIM**
Nama: Pensyarah
Cop Rasmii: Jabatan Sains Biologi
Kolej Universiti Sains dan Teknologi Malaysia
(KUSTEM)
21030 Kuala Terengganu, Terengganu.

Tarikh: 14/65/06

Ketua Jabatan Sains Biologi

Nama: **PROF. MADYA DR. NAKISAH BT. MAT AMIN**
Cop Rasmii: Ketua
Jabatan Sains Biologi
Fakulti Sains dan Teknologi
Kolej Universiti Sains dan Teknologi Malaysia
(KUSTEM)
21030 Kuala Terengganu.

Tarikh: 21.5.06

ACKNOWLEDGEMENTS

Alhamdulillah. With the name of Allah, the most gracious and merciful.

I would like express my deepest gratitude to my supervisor, Cik Jamilah Bt. Mohd Salim. Thank you for the support, caring and patient supervision throughout this project.

My thank also goes to En. Razali Salam for the valuable guidance and helps and all staff of FRIM research station in Setiu, Terengganu for their assistance.

I also would like to thanks my parents, Sabri and Roslah Abd Ghani, my brother and sisters, Kakngah, Anis and Fahmi, for their love, courage and support. To my beloved friends and housemates, Yan(Nek), Atie, Ala, Fara, Sal, Ira, Aza, Mok Ani, Kak Rumie, Kak Pujah, Kak Zila, McLah and Wani. Thank you for being with me and sharing.

Lastly, thank you to those names that I might have omitted, who ever have helped to make this project possible.

TABLE OF CONTENTS

Title	Page
ACKNOWLEDGEMENTS	ii
TABLE OF CONTENTS	iii
LIST OF TABLES	vi
LIST OF FIGURES	v
LIST OF ABBREVIATIONS	vi
LIST OF APPENDICES	viii
ABSTRACT	ix
ABSTRAK	x

CHAPTER 1 INTRODUCTION

1.1 Introduction	1
1.2 Objectives	3

CHAPTER 2 LITERATURE REVIEW

2.1 <i>Ficus</i>	4
2.2 Ecosystem of BRIS Soil	5
2.3 Seedling growth	6

CHAPTER 3 METHODOLOGY

3.1 Site Description	9
3.2 Seedlings growth in natural habitat	10
3.3 Seedlings growth in artificial medium	10
3.4 Data analysis	12

CHAPTER 4 RESULT

4.1 Site Description	13
4.2 Seedlings growth in natural habitat	14
4.3 Seedlings growth in artificial medium	15

CHAPTER 5 DISCUSSION

5.1 Seedlings growth in natural habitat	17
5.1 Seedling growth in artificial medium	17

CHAPTER 6 CONCLUSION

REFERENCES	19
-------------------	----

APPENDICES	20
-------------------	----

CURICULUM VITAE	23
------------------------	----

28

LIST OF TABLES

Table	Page
4.1 Descriptive statistics from seedlings height and diameter of <i>F.deltoidea</i> in natural habitat of BRIS soil of Bari, Setiu and Marang Terengganu.of height plant for first time recorded in Bari and Marang.	14
4.2 Descriptive statistic of light intensity under 20 clumps of <i>R.tomentosa</i> that occupied by <i>F.deltoidea</i> in Setiu and Marang	14

LIST OF FIGURES

Figure		Page
1.1	<i>F.deltoidea</i> in its natural habitat under the clump of <i>Rhodomyrtus Tomentosa</i>	2
3.1	Sampling location at Kg. Bari Besar, Setiu and Kg. Pulau Kerengga, Marang.	9
3.2	Seeds arranged systematically in tray with toothpick to easily spot the emergence of plumule.	11
3.3	Methods adopted to germinate seeds of <i>F.deltoidea</i> .	11
4.1	The study area on heath vegetation with BRIS soil in Setiu, Terengganu.	13
4.2	The condition <i>F. deltoidea</i> clump after being collected by the locals in Setiu, Terengganu.	15
4.3	Germination percentage of <i>F.deltoidea</i> seeds in three medium of BRIS soil, sand and litter.	16

LIST OF ABBREVIATIONS

cm	centimetre
Kg	Kampung
N	number of samples
%	Percentage
"	Minutes
°	Degree
BRIS	Beach Ridges Interspersed with Swales

ABSTRACT

Seedlings growth of *Ficus deltoidea* Jack. in natural habitat and artificial medium was compared. Germination on BRIS soil medium was better than sand or litter medium. The result of this study has shown that the germination percentage of *F.deltoidea* in BRIS soil medium was 90%. Shoots were produced after seven days of cultivation. Mean height on BRIS soil medium was 1.93 ± 0.06 cm and 4.90 ± 0.07 mean number of leaf after ten weeks. Mean height on litter medium after ten weeks was 4.73 ± 0.07 and 4.90 ± 0.07 leaves number. The extreme light intensity on BRIS soil may limit the regeneration of *F.deltoidea* seedling outside *Rhodomyrtus tomentosa* clumps in natural habitat on BRIS soil ecosystem.

PERBANDINGAN PERTUMBUHAN ANAK BENIH *FICUS DELTIDEA* JACK. DIANTRA HABITAT SEMULAJADI DAN MEDIUM TANAH

ABSTRAK

Perbandingan pertumbuhan anak benih *Ficus deltoidea* Jack. telah dilihat diantara habitat semulajadi dan menggunakan medium tanah. Percambahan menggunakan medium tanah BRIS adalah lebih baik daripada pasir atau sarap. Hasil kajian menunjukkan peratusan percambahan *F.deltoidea* di tanah BRIS adalah 90%. Pertumbuhan pucuk mula kelihatan selepas tujuh hari dalam semaian. Purata ketinggian pokok di medium tanah BRIS ialah 1.93 ± 0.06 cm dan purata bilangan daun 4.90 ± 0.07 helai selepas sepuluh minggu. Purata tinggi pokok dalam medium sarap adalah 4.73 ± 0.07 cm dengan purata 4.90 ± 0.07 helai daun. Keamatan cahaya melampau di tanah BRIS mungkin menghadkan regenerasi anak benih *F.deltoidea* di luar kelompok *Rhodomyrtus tomentosa* di habitat semulajadi di atas ekosistem tanah BRIS.