

A STUDY ON THE GROWTH RATE  
BETWEEN NATURAL AND TRANSPLANTED SEEDLINGS OF  
*Rhizophora apiculata* BL.

FARIDA HANIM BT ABDUL RAHIM

FACULTY OF APPLIED SCIENCE AND TECHNOLOGY  
UNIVERSITI PUTRA MALAYSIA  
TERENGGANU

1999

LP  
10  
FSGT  
2  
1999

1100024124

PERPUSTAKAAN  
UNIVERSITI PUTRA MALAYSIA TERENGGANU

LP 10 FSGT 2 1999



1100024124

## A study on the growth rate between natural and transplanted seedlings of *Rhizophora apiculata* BL / Faridah Hanim Abdul Rahim.

**PERPUSTAKAAN**  
**KOLEJ UNIVERSITI SAINS & TEKNOLOGI MALAYSIA**  
**21030 KUALA TERENGGANU**

1100024124

1100024124

Lihat sebelah

HAK MILIK  
PERPUSTAKAAN KUSTEM

A STUDY ON THE GROWTH RATE  
BETWEEN NATURAL AND TRANSPLANTED SEEDLINGS OF

*Rhizophora apiculata* BL.

BY

FARIDA HANIM BT ABDUL RAHIM

This project report is submitted in partial fulfillment of

the requirements for

the Degree of Bachelor of Marine Science

Faculty of Applied Science and Technology

UNIVERSITI PUTRA MALAYSIA TERENGGANU

1999

1100024124

## **ACKNOWLEDGEMENT**

A great many people have contributed to the production of this thesis. For substantial financial assistance, I wish to thank University Putra Malaysia Terengganu (UPMT), particularly Mr. Sulong Ibrahim as my first supervisor. I also owe a debt of gratitude to Prof. Madya Dr. Lokman Hussien as my second supervisor for their support and sponsorship.

I wish to express my sincere thanks to my beloved one; Anuar Yusof and my family for their encouragement and moral support for me to finish my project successfully.

Many thanks to Mr. Abdull Habir, Mohd Tarmizi, Mohd Harith, Mohd Razali Salam, Mohd Suffian, Abdul Razak and Helini who contributed their time and knowledge. I am most grateful for their companionship in circumstances that were not always particularly comfortable.

Similarly, I am fully indebted to all my friends, Safiyya , Abdul Wahab, Zool Fadly, Mohd Anuar and Zailani, who worked with me in most co-operative, collegial and friendly way on the finishing of my project.

Last but not least, I wish to express my deep gratitude to all those people, named and unnamed, who have supported and encouraged me through my years of preparing this thesis, I offer my heartfelt thanks.

## ABSTRAK

Kajian perbandingan tumbesaran antara anak pokok semulajadi dan anak pokok yang ditanam semula belum dijalankan sepenuhnya. Dua ratus anak pokok *Rhizophora apiculata* BL telah dibawa dari kawasan semulajadinya di Bakau Tinggi, Kemaman ke UPMT untuk ditanam semula bagi kajian perbandingan kadar tumbesaran dan kadar kemandirian. *Rhizophora apiculata* BL dipilih untuk kajian ini kerana ia mempunyai ciri-ciri bakau yang jelas kelihatan seperti akarnya yang unik dan kehadirannya yang sedikit di UPMT. Data-data ketinggian, ukur lilit dan bilangan daun bagi setiap anak pokok dikutip setiap dua bulan. Kadar kemandirian bagi anak pokok yang ditanam semula adalah 91.25%. Walaupun t-test menunjukkan bahawa anak pokok semulajadi tumbuh dengan lebih cepat berbanding anak pokok yang ditanam semula, kadar 65% tumbesaran berbanding anak pokok semulajadi menunjukkan tumbesaran kedua-dua anak pokok adalah sama bagi jangkamasa yang tertentu. Kenaikan purata isipadu anak pokok di Bakau Tinggi ialah  $4.25\text{ cm}^3$  manakala di UPMT ialah  $2.76\text{ cm}^3$ . Purata kehilangan daun bagi setiap anak pokok di Bakau Tinggi ialah 1.65 helai manakala di UPMT ialah 3.86 helai. Kematian anak pokok yang ditanam semula di UPMT mungkin disebabkan oleh tekanan yang dihadapi ketika proses penanaman semula dijalankan. Diketahui bahawa sekitaran anak pokok yang ditanam semula adalah berbeza dengan sekitaran semulajadinya. Sekitaran anak pokok yang ditanam semula di UPMT adalah lebih berpasir dan ditenggelami air ketika musim hujan. Ketika musim tengkujuh, buaan di mulut muara selalu tertutup dan menyebabkan air hujan dan enapan terperangkap di kebanyakan kawasan sekitar UPMT. Ini membuktikan bahawa penanaman semula di kawasan yang kurang baik keadaan sekitarannya masih boleh diusahakan dengan jayanya.

## ABSTRACT

A comparative study of growth between natural and transplanted seedlings is not well documented. Two hundred seedlings of *Rhizophora apiculata* BL were transferred from its natural area at Kemaman to UPMT in an effort to compare their growth and survivability rate. *Rhizophora apiculata* BL was chosen for this study because of their presence is not lush and they have unique flying buttress roots. Data of height, circumference and numbers of leaves were collected from each location; UPMT and Bakau Tinggi. The survivability rate of the transplanted seedlings is 91.25% within four month. Although t-test revealed that the growth rate of natural seedlings is faster than the transplanted seedlings, the fact that the transplanted seedlings could grow at 65% the rate of natural seedlings, indicates that given time the growth rate between the two groups of seedlings could be similar. The monthly increment of average volume at Bakau Tinggi is  $4.25 \text{ cm}^3$  while at UPMT is  $2.76 \text{ cm}^3$ . The average numbers of leaves lost at Bakau Tinggi is 1.65 while at UPMT is 3.86. It was found that the transplanted condition is quite different from the natural condition. It is more sandy and not well flushed. The sea water enters UPMT through an opening at the beach. During monsoon season, the opening was closed and trapped the rainwater and sediments around UPMT. Whole seedlings were drowned and the sediments covered their leaves. The seedlings also shocked during transplanting process, which affected their growth. It can be concluded that, at least for *Rhizophora apiculata* BL species, the effort of transplanting to a non-hospitable condition could be successfully done.