

AIDA ROYYANI BINTI ABD ARIFF

MASTER OF SCIENCE

2015

**DISTRIBUTION OF WATER PHYSICO-
CHEMICAL PARAMETERS AND BIOGENIC
ELEMENTS IN SETIU LAGOON DURING
MONSOON EVENTS AS INDICATOR FOR
PRIMARY PRODUCTIVITY**

AIDA ROYYANI BINTI ABD ARIFF

**MASTER OF SCIENCE
UNIVERSITI MALAYSIA TERENGGANU**

2015

**DISTRIBUTION OF WATER PHYSICO-CHEMICAL PARAMETERS AND
BIOGENIC ELEMENTS IN SETIU LAGOON DURING MONSOON
EVENTS AS INDICATOR FOR PRIMARY PRODUCTIVITY**

AIDA ROYYANI BINTI ABD ARIFF

**Thesis Submitted in Fulfillment of the Requirement for the
Degree of Master of Science in the School of Marine and
Environmental Sciences
Universiti Malaysia Terengganu**

June 2015



This thesis was affectionately dedicated to
my family members...

Especially to my mum and dad,
who always encouraged me
to go on every difficulties...
their endless love always clinging along
my pathway of success...

I love you all...



Abstract of thesis presented to the Senate of Universiti Malaysia
Terengganu in fulfillment of the requirement for the degree of Master of
Science

**DISTRIBUTION OF WATER PHYSICO-CHEMICAL PARAMETERS AND
BIOGENIC ELEMENTS IN SETIU LAGOON DURING MONSOON
EVENTS AS INDICATOR FOR PRIMARY PRODUCTIVITY**

AIDA ROYYANI BINTI ABD ARIFF

June 2015

Main Supervisor : Zainuddin bin Bachok, Ph.D.
**Co-Supervisor : Assoc. Prof. Mohamed Kamil bin Abdul Rashid,
Ph.D.**
School : School of Marine and Environmental Sciences

The study was conducted in the Setiu Lagoon, Terengganu involving three sampling periods, namely in May 2007 (post-monsoon), September 2007 (pre-monsoon) and January 2008 (monsoon). Samples of nutrients (NO_3^- -N, NO_2^- -N, NH_4^+ -N, PO_4^{3-} -P), primary productivity, chlorophyll-a and physico-chemical parameters of water were collected during low and high tides from 6 stations in the lagoon area and 6 stations in the riverine area except for primary productivity where samples were taken during the pre-monsoon and monsoon seasons only during high tide. Data obtained were analyzed to assess the variations of water quality within Setiu lagoon spatially and temporally subjected to monsoonal events and tidal changes. From the results, the N:P ratio indicated that the study area was mostly N limited especially in the lagoon area. Correlation analysis indicated that phytoplankton consumed more NO_3^- -N during low tide in the post-monsoon ($p < 0.01$) and pre-monsoon season ($p < 0.05$) but during high tide,

$\text{PO}_4^{3-}\text{-P}$ more favorable to phytoplankton in the post-monsoon ($p < 0.01$) and pre-monsoon season ($p < 0.01$). Factor analysis showed that monsoonal event affects the phytoplankton productivity in the study area compared to tidal event. Temperature was the primary factor while nutrients concentration and salinity were the secondary factors. These indicated that monsoonal events coupled with tidal events contribute to the nutrient supply into the study area. Non-point sources and natural sources were the contributor for the nutrient enhancement in the study area. The study area was still not polluted with nutrients loading and did not exhibit any sign of eutrophication. The nutrient level did not exceed the standard of MWQCS (2010). The physico-chemical parameters and nutrients can be a good indicator for primary productivity in the Setiu lagoon.

Abstrak tesis yang dikemukakan kepada Senat Universiti Malaysia Terengganu sebagai memenuhi keperluan untuk Ijazah Master Sains

TABURAN PARAMETER FISIKO-KIMIA AIR DAN UNSUR BIOGENIK DI LAGUN SETIU SEMASA MONSOON SEBAGAI PENUNJUK PRODUKTIVITI PRIMER

AIDA ROYYANI BINTI ABD ARIFF

Jun 2015

Penyelia Utama : Zainuddin bin Bachok, Ph.D.

Penyelia Bersama : Assoc. Prof. Mohamed Kamil bin Abdul Rashid, Ph.D.

Pusat Pengajian : Pusat Pengajian Sains Marin dan Sekitaran

Kajian ini telah dijalankan di Lagun Setiu, Terengganu melibatkan tiga kali persampelan iaitu pada Mei 2007 (post-monsoon), September 2007 (pra-monsoon) dan Januari 2008 (monsoon). Sampel nutrien (NO_3^- -N, NO_2^- -N, NH_4^+ -N, PO_4^{3-} -P), produktiviti primer, klorofil-a, dan parameter fisiko-kimia air diambil ketika air surut dan air pasang dari 6 stesen di kawasan lagun and 6 stesen di kawasan sungai kecuali untuk sampel produktiviti primer yang diambil hanya ketika musim pra-monsoon dan monsoon ketika air pasang. Data yang diperolehi dianalisa untuk melihat variasi kualiti air di lagun Setiu mengikut kawasan dan masa berdasarkan acara monsoon dan perubahan pasang-surut. Dari keputusan yang diperolehi, nisbah N:P menunjukkan kawasan kajian adalah dihadkan oleh N terutamanya di kawasan lagun. Analisis korelasi menunjukkan fitoplankton menggunakan NO_3^- -N semasa air surut pada musim post-monsoon ($p < 0.01$) dan pra-monsoon ($p < 0.05$) tetapi semasa air pasang, PO_4^{3-} -P lebih digemari oleh fitoplankton pada musim post-monsoon ($p < 0.01$) and pra-monsoon ($p < 0.01$).

Analisis factor menunjukkan bahawa acara monsun lebih memberi kesan terhadap produktiviti fitoplankton di kawasan kajian berbanding dengan acara pasang-surut. Suhu merupakan faktor primer manakala kepekatan nutrien dan kemasinan merupakan factor sekunder. Ini menunjukkan bahawa acara monsun bersama dengan acara pasang-surut berperanan di dalam membawa bekalan nutrien ke dalam kawasan kajian. Sumber bukan titik dan sumber semulajadi adalah penyumbang kepada peningkatan nutrien di kawasan kajian. Kawasan kajian masih lagi tidak tercemar dengan kemasukan nutrien dan tidak menunjukkan sebarang tanda eutrofikasi. Tahap nutrien masih di bawah atau tidak melebihi piawaian yang telah ditetapkan oleh MWQCS (2010). Parameter fisiko-kimia dan nutrien boleh dijadikan sebagai penunjuk yang baik kepada produktiviti primer di lagun Setiu.