

NK KELLOGGIANA BUDWELLER
BLOOMING IN THE RECENT STAGE
OF BLOOMING INDICATED
THEY ARE IN THE ACTIVE STAGE

DALE KELLOGG

KELLOGG COMPANY CHIEF BOTANIST
KELLOGG UNIVERSITY CHIEF BOTANIST MALAYSIA
2000

CM: 4734

1100046041 Perpustakaan
Universiti Malaysia Terengganu (UMT)

LP 40 FST 3 2006



1100046041

Efficacy of tank mixing ametryn with glyphosate or glufosinate-ammonium on growth of goosegrass (*Eleusine Indica* (L) gaertn) at the mature stage / Nor Asmah Jalil.




PERPUSTAKAAN
KOLEJ UNIVERSITI SAINS & TEKNOLOGI MALAYSIA
21030 KUALA TERENGGANU

1100046041

1100046041

Lihat sebelah



**EFFICACY OF TANK MIXING AMETRYN WITH GLYPHOSATE OR
GLUFOSINATE-AMMONIUM ON GROWTH OF GOOSEGRASS
(*ELEUSINE INDICA* (L.) GAERTN.) AT THE MATURE STAGE**

NOR ASMAH BINTI JALIL

**FAKULTI SAINS DAN TEKNOLOGI
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MALAYSIA
2006**

EFFICACY OF TANK MIXING AMETRYN WITH GLYPHOSATE OR
GLUFOSINATE-AMMONIUM ON GROWTH OF GOOSEGRASS
(ELEUSINE INDICA (L.) GAERTN.) AT THE MATURE STAGE

By

Nor Asmah binti Jalil

Research Report submitted in partial fulfillment of
the requirements for the degree of
Bachelor of Science (Biological Sciences)

Department of Biological Sciences
Faculty of Science and Technology
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MALAYSIA
2006

This project should be cited as:

Asmah, J.N. 2006. Efficacy of tank mixing ametryn with glyphosate or glufosinate-ammonium on growth of goosegrass (*Eleusine indica* (L.) Gaertn) at the mature stage. Undergraduate thesis. Bachelor of Science in Biological Sciences, Faculty of Science and Technology, Kolej Universiti Sains dan Teknologi Malaysia, Terengganu. 60p.

No part of this project report may be produced by any mechanical, photographic, or electronic process, or in the form of phonographic recording, nor may it be stored in a retrieval system, transmitted, or otherwise copied for public or private use, without written permission from the author and supervisor(s) of the project.



JABATAN SAINS BIOLOGI
FAKULTI SAINS DAN TEKNOLOGI
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MALAYSIA

PENGAKUAN DAN PENGESAHAN LAPORAN
PROJEK PENYELIDIKAN I DAN II

Adalah ini diakui dan disahkan bahawa laporan penyelidikan bertajuk: EFFICACY OF TANK MIXING AMETRYN WITH GLYPHOSATE OR GLUFOSINATE-AMMONIUM ON GROWTH OF GOOSEGRASS (*Eleusine indica* (L.) Gaertn.) AT THE MATURE STAGE oleh Nor Asmah binti Jalil, no. matrik: UK 8576 telah diperiksa dan semua pembetulan yang disarankan telah dilakukan. Laporan ini dikemukakan kepada Jabatan Sains Biologi sebagai memenuhi sebahagian daripada keperluan memperolehi Ijazah Sarjana Muda Sains (Sains Biologi) Fakulti Sains dan Teknologi, Kolej Universiti Sains dan Teknologi Malaysia.

Disahkan oleh:

.....
Penyelia Utama

Nama: Dr. Cha Thye San
Kolej Universiti Sains dan Teknologi Malaysia
Cop Rasmi:
DR. CHA THYE SAN
Pensyarah
Jabatan Sains Biologi
Fakulti Sains dan Teknologi
(KUSTEM)
21030 Kuala Terengganu.

Tarikh: 27/4/2006

.....
Penyelia Kedua (jika ada)

Nama: Dr. Chuah Tse Seng

Cop Rasmi

Dr. Chuah Tse Seng
Pensyarah
Jabatan Agroteknologi
Fakulti Agroteknologi dan Sains Makanan
Kolej Universiti Sains dan Teknologi Malaysia
21030 Kuala Terengganu.

.....
Tarikh: 4/5/2006

.....
Ketua Jabatan Sains Biologi

Nama: Prof. Madya Dr. Nakisah Bt. Mat Amin

Cop Rasmi:
PROF. MADYA DR. NAKISAH BT. MAT AMIN
Kezia
Jabatan Sains Biologi
Fakulti Sains dan Teknologi
Kolej Universiti Sains dan Teknologi Malaysia
(KUSTEM)
21030 Kuala Terengganu.

.....
Tarikh: 27/4/06

ACKNOWLEDGEMENT

A special thank goes to Dr. Chuah Tse Seng for many hours of work, suggestion, tremendous patience and encouragement. He thought me how to express my idea and showed me a different ways to approach a research problem. I'm also thank my supervisor, Dr. Cha Thye San who always undying encouragement. Sincere thank also goes to En. Hassan who has been willing to help during conducted study in the greenhouse.

Let me also say 'thank you' to my friends, Shirley Maclaine, Franci Julop, Anne Marie Caben, Suganthi Krishnan, Nor Shasha Zura Md. Lajis, Mazlina, Wan Azizan Abdul Aziz, and Siti Nurul Husna for their friendship, many hours of work and sweet memories. I'm also thank to my lovely family who give support, motivation, listening to my complaints and frustration and their contribution to the overall appearance to finish this research.

And last but not least, I want to express thank you to all the people who involved and gave a lot of encouragement in supporting this research. Their helpful criticisms and suggestion for improvement were significant contributions that I'm greatly appreciated. Thank you.

CONTENTS

	Page
ACKNOWLEDGEMENT	ii
LIST OF TABLES	vi
LIST OF FIGURES	vii
LIST OF ABBREVIATIONS / SYMBOLS	viii
LIST OF APPENDICES	ix
ABSTRACT	x
ABSTRAK	xi
CHAPTER 1 INTRODUCTION	1
CHAPTER 2 LITERATURE REVIEW	
2.1 Weeds	5
2.2 Eleusine indica (L.) Gaertn	6
2.2.1 Habitat and ecology	6
2.2.2 Distribution	7
2.2.3 Morphology	7
2.2.4 Agricultural imparatnce	10
2.3 Chemical weeds control	11
2.3.1 Herbicide classification	11
2.3.2 Adoption of herbicides	11
2.4 Herbicides	13

2.4.1	Triazine herbicides	13
2.4.1a	Ametryn	14
2.4.2	Glyphosate	16
2.4.3	Glufosinate-ammonium	18
2.5	Tank-mixing	20
2.5.1	Herbicide joint action	22

CHAPTER 3 METHODOLOGY

3.1	Plants materials	24
3.2	Herbicides	25
3.3	Dose-response experiment	25
3.4	Herbicide combination experiment	28
3.5	Field experiment	31

CHAPTER 4 RESULTS

4.1	Dose response of Eleusine indica (L.) Gaertn. to different rates of herbicides	32
4.1.1	Dose response of Eleusine indica to ametryn	32
4.1.2	Dose response of Eleusine indica to glyphosate	34
4.2	Combination of glyphosate plus ametryn on goosegrass control	36
4.3	Combination of glufosinate-ammonium plus ametryn on goosegrass control	38
4.4	Visual assessment on herbicides combination in the fields	40

CHAPTER 5	DISCUSSION	42
CHAPTER 6	CONCLUSION	47
REFERENCES		48
APPENDICES		52

LIST OF TABLES

Tables		page
3.1	Herbicides rates for dose-response experiment at the mature stage	26
3.2	Herbicides rates for herbicides combination experiments	30
4.1	Goosegrass shoot weight reduction in the greenhouse four weeks after treated with combinations of glyphosate plus ametryn	37
4.2	Goosegrass shoot weight reduction in the greenhouse four weeks after treated with combinations of glufosinate-ammonium plus ametryn	39
4.3	Percentage of visual assessment for goosegrass control at four and six Weeks after tank mixture treatment	41

LIST OF FIGURES

Figures

pages

2.1	<i>Eleusine indica</i> (L.) Gaertn	9
2.2	Molecular structure of ametryn	14
4.1	Log logistic response regression line of goosegrass (<i>Eleusine indica</i> (L.) Gaertn.) in different rates of ametryn	33
4.2	Log logistic response regression line of goosegrass (<i>Eleusine indica</i> (L.) Gaertn.) in different rates of glyphosate	35

LIST OF ABBREVIATIONS / SYMBOLS

cm	=	Centimeter
g	=	Gram
kg	=	Kilogram
a. i. ha ⁻¹	=	Active ingredient per hectare
a. i. L ⁻¹	=	Active ingredient per Liter

LIST OF APPENDICES

Appendices		Page
A	Regression analysis of <i>Eleusine indica</i> (L.) Gaertn. , after application of ametryn	53
B	Regression analysis of <i>Eleusine indica</i> (L.) Gaertn. , after application of glyphosate	54
C	Responses of <i>Eleusine indica</i> (L.) Gaertn to glyphosate at different rates	55
D	Responses of <i>Eleusine indica</i> (L.) Gaertn to ametryn at different rates	56
E	Comparison cost of single herbicide and herbicide mixtures (glufosinate-ammonium plus ametryn and glyphosate plus ametryn) on goosegrass shoot weight reduction in three weeks after herbicide treatment.	57

ABSTRACT

A study was conducted to determine the efficacy of tank mixtures of ametryn plus glyphosate or glufosinate-ammonium on control of goosegrass at the mature stage in the greenhouse. Fresh weight reduction of plants treated with glyphosate plus ametryn mostly were more than predicted value of fresh weight reduction indicating additive responses between these two herbicides. Six of nine tank mixtures were additive while three of them were antagonistic. Shoot fresh weight reduction of plants treated with glufosinate-ammonium plus ametryn were also showed that mostly were additive responses of these combination. Additive response was observed in five tank mixtures while other four tank mixtures gave antagonistic response. Good control of goosegrass applied in the field is in the line with the result obtained in the greenhouse study. The most cost effective control of herbicide combination treatment is to mix 320 g a. i. ha⁻¹ glyphosate with 80 g a. i. ha⁻¹ ametryn.

**KEBERKESANAN KOMBINASI AMETRIN DAN GLIFOSAT ATAU
GLUFOSINAT-AMMONIUM UNTUK KAWALAN RUMPUT SAMBARI
(*Eleusine indica* (L.) Gaertn) PADA PERINGKAT MATANG.**

ABSTRAK

Kajian ini telah dijalankan mengkaji keberkesanannya kombinasi glifosat dan ametrin serta kombinasi glufosinat-ammonium dan ametrin untuk kawalan rumput sambari pada peringkat matang di rumah hijau. Pengurangan berat basah bagi semua kombinasi glifosat dan ametrin menunjukkan kebanyakannya adalah lebih tinggi berbanding dengan nilai pengurangan berat basah ramalan, menunjukkan bersifat aditif. Enam daripadanya adalah aditif manakala tiga campuran lagi adalah antagonistik. Pengurangan berat basah bagi rumput sambari yang di uji dengan kombinasi glufosinat-ammonium dan ametrin juga menunjukkan kebanyakannya adalah bersifat aditif. Lima daripada sembilan kombinasi adalah aditif manakala empat lagi adalah antagonistik. Kombinasi kawalan yang baik terhadap rumput sambari yang disembur di lapangan juga menunjukkan keputusan yang sama dengan keputusan kombinasi kawalan di rumah hijau. Kombinasi yang paling baik adalah kombinasi 320 g a. i. ha^{-1} glifosat dan 80 g a. i. ha^{-1} ametrin.